



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

it with *Emberiza* Brisson, but an examination of its type and comparison with typical species of *Emberiza* shows that it is well differentiated as a generic group. It differs from *Emberiza* Brisson (type, by tautonymy, *Emberiza citrinella* Linnaeus) as follows; bill slenderer, more compressed, more sharply pointed, thus less conical; basal two-thirds of culmen straight or even somewhat concave, instead of convex; maxillar and mandibular tomia vertically not so strongly concave, thus not giving the closed commissure the somewhat open appearance that it has in typical species of *Emberiza*; palatal surface of maxilla lacking the peculiar rounded protuberances of *Emberiza*; mandible more rounded (less squarish) basally; gonys very long, its length much more than the height of the bill at base (instead of about equal to that dimension), and not strongly ascending, the gonydeal angle therefore not so prominent; tertials and tail much shorter.

The species to be included in this genus are at least the three originally indicated by Cabanis, the last one of which is North American by reason of its accidental occurrence on Kiska Island in the Aleutian Islands, Alaska. These are:

Hypocentor aureolus (Pallas).

Hypocentor fucatus (Pallas).

Hypocentor rusticus (Pallas).

HARRY C. OBERHOLSER, Washington, D. C.

A Correction Involving Some Juncos.—An error that may be explained as due to oversight, inadvertence, plain stupidity or all three combined, crept into my paper on the Juncos (Bull. Am. Mus. Nat. Hist. XXXVIII, 1918, p. 296) and Mr. Todd has called my attention to it. In placing *insularis* under *mearnsi* as a race, I quite forgot that the former name has many years priority. Therefore the Pink-sided Juncos should stand as follows:—

Junco insularis mearnsi

Junco insularis insularis

Junco insularis townsendi

JONATHAN DWIGHT, M. D. New York City.

An Additional Record of *Ammodramus savannarum bimaculatus* in Eastern Washington.—Although the breeding range of the Western Grasshopper Sparrow is stated by the Check List (A. O. U. Check-List of North American Birds, 1910, p. 257) to embrace "Transition and Austral zones from southeastern British Columbia, northwestern Montana, and southern Minnesota south to southern California and southern Texas," it appears that only one actual record of occurrence in eastern Washington has been published to date. Dr. Lee R. Dice took two adult males in breeding plumage in a wheat field in the Touchet Valley, near Prescott, Walla Walla County, on June 16, 1908 (Auk, Vol. XXVII, 1910, p. 217).